

RHODE ISLAND

Table 89 Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Rhode Island, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2011-2012 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	140	12	44	84	128
Past Year Marijuana Use	174	16	61	98	159
Past Month Marijuana Use	116	10	40	67	107
Past Month Use of Illicit Drugs Other Than Marijuana ¹	42	3	13	26	39
Past Year Cocaine Use	23	1	9	14	23
Past Year Nonmedical Pain Reliever Use	47	4	15	27	43
Perception of Great Risk of Smoking Marijuana Once a Month	214	16	15	183	198
Average Annual Number of Marijuana Initiates ²	12	5	6	1	7
ALCOHOL					
Past Month Alcohol Use	567	12	97	457	554
Past Month Binge Alcohol Use ³	269	7	71	191	263
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	334	29	40	265	305
Past Month Alcohol Use (Persons Aged 12 to 20)	40 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	26 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	251	8	54	190	243
Past Month Cigarette Use	194	6	44	144	188
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	649	51	89	508	598
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	21	2	8	11	20
Illicit Drug Dependence or Abuse ¹	27	4	11	13	24
Alcohol Dependence	35	1	12	22	34
Alcohol Dependence or Abuse	80	3	25	51	77
Alcohol or Illicit Drug Dependence or Abuse ¹	95	5	31	59	89
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	23	3	9	10	20
Needing But Not Receiving Treatment for Alcohol Use ⁷	74	3	25	47	71
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	7	12	51	62
Serious Mental Illness ^{9,10}	--	--	6	29	35
Any Mental Illness ^{9,11}	--	--	25	129	154
Had Serious Thoughts of Suicide	--	--	9	24	33

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "12+" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older.

⁹ For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <http://www.samhsa.gov/data/NSDUH/2k12State/NSDUHsae2012/Index.aspx>.

¹⁰ Serious mental illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment.

¹¹ Any mental illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 (SMI and AMI Estimates Revised October 2013) and 2012.

RHODE ISLAND

Table 90 Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Rhode Island, by Age Group: Percentages, Annual Averages Based on 2011-2012 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ¹	15.61	15.16	32.88	12.33	15.66
Past Year Marijuana Use	19.47	20.11	46.03	14.26	19.41
Past Month Marijuana Use	13.00	12.44	30.16	9.74	13.05
Past Month Use of Illicit Drugs Other Than Marijuana ¹	4.70	4.26	9.60	3.81	4.75
Past Year Cocaine Use	2.62	0.80	6.82	2.02	2.79
Past Year Nonmedical Pain Reliever Use	5.23	5.46	11.66	3.96	5.21
Perception of Great Risk of Smoking Marijuana Once a Month	23.89	20.08	11.39	26.76	24.25
Average Annual Rate of First Use of Marijuana ²	2.74	8.61	10.94	0.26	1.75
ALCOHOL					
Past Month Alcohol Use	63.35	15.74	73.46	66.80	67.88
Past Month Binge Alcohol Use ³	30.11	8.63	53.92	27.94	32.15
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	37.28	37.03	30.28	38.66	37.30
Past Month Alcohol Use (Persons Aged 12 to 20)	31.48 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	20.44 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	28.05	9.97	40.49	27.70	29.77
Past Month Cigarette Use	21.70	7.72	33.33	21.05	23.03
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	72.51	65.64	67.23	74.31	73.17
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	2.39	2.43	6.26	1.64	2.39
Illicit Drug Dependence or Abuse ¹	3.07	4.89	8.11	1.89	2.89
Alcohol Dependence	3.87	1.40	8.77	3.20	4.10
Alcohol Dependence or Abuse	8.92	3.87	19.22	7.51	9.40
Alcohol or Illicit Drug Dependence or Abuse ¹	10.57	6.89	23.11	8.56	10.91
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.56	4.05	7.06	1.52	2.42
Needing But Not Receiving Treatment for Alcohol Use ⁷	8.30	3.72	18.75	6.80	8.74
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	9.00	8.79	7.43	7.65
Serious Mental Illness ^{9,10}	--	--	4.27	4.25	4.25
Any Mental Illness ^{9,11}	--	--	18.80	18.81	18.80
Had Serious Thoughts of Suicide	--	--	6.89	3.51	4.05

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual marijuana initiation rate = $100 * \{ [X_1 \div (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "12+" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older.

⁹ For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <http://www.samhsa.gov/data/NSDUH/2k12State/NSDUHsae2012/Index.aspx>.

¹⁰ Serious mental illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment.

¹¹ Any mental illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 (SMI and AMI Estimates Revised October 2013) and 2012.